



GRAPHIC ILLUSTRATIONS: PHARMACODYNAMICS & PHARMACOKINETICS**Time Course of Drug Action**

- onset of action: the time, after a drug is administered, to achieve a drug serum concentration required to produce a detectable response ($t_0 \rightarrow t_1$)
- time to peak: the time required for a drug to achieve its highest therapeutic serum concentration ($t_0 \rightarrow t_2$)
- duration of action (DOA): the amount of time a drug is present in adequate serum concentration necessary to produce a therapeutic effect ($t_1 \rightarrow t_3$)
 - DOA depends on the rate of drug absorption and elimination
- half-life ($t_{1/2}$): the amount of time required for elimination processes to reduce the drug serum concentration by one-half



Continuous vs Intermittent Drug Administration



Estimated Creatinine Clearance (for drug dosing considerations)

A. Calculation of Ideal Body Weight (IBW)

1. Female

$$\text{IBW} = (2.3 \times \text{inches above 5'}) + 45.5 \text{ kg}$$

2. Male

$$\text{IBW} = (2.3 \times \text{inches above 5'}) + 50 \text{ kg}$$

B. Creatinine Clearance (CrCl) Formula

$$\text{CrCl (ml/min)} = \frac{(140 - \text{Age}) \times \text{IBW}}{\text{sCr} \times 72}$$

(sCr = serum creatinine level)

$$\text{CrCl}_{\text{female}} = \text{CrCl} \times 0.85$$

C. Adjusted Body Weight (ABW)

$$\text{ABW} = \text{IBW} + 0.4 (\text{Actual Body Wt} - \text{IBW})$$

CREATININE CLEARANCE SAMPLE PROBLEM

NOTE: The CrCl formula will be provided to students during the exam. The emphasis will be on application and making the right assessment.

Jane Smith (JS) is a 69-year old female who was admitted to University Hospital complaining of UTI symptoms. Dr. Urobact, Jane's physician, would like you to recommend an appropriate dose of Keflex (cephalexin), based on Jane's kidney function.

Jane's HT / WT = 5' 4" / 152 LBS.

A blood chemistry panel reported the following values:

Na = 145
K = 3.8
Cl = 101
Glucose = 98
CO₂ = 24
BUN = 27
Cr = 1.5

QUESTIONS

1. Calculate Jane's IBW (ideal body wt)
2. Calculate her CrCl (creatinine clearance).
3. Recommend the appropriate dose of Keflex for Jane.

KEFLEX (Cephalexin) Package Insert

According to the package insert of Keflex (cephalexin), 500 mg every 6 hours is recommended for a CrCl > 50 ml/min. But, if the patient's kidney function reflects a range of 10 to 50 ml/min, the dose of cephalexin should be reduced to 500 mg every 8 to 12 hours. For CrCl < 10 ml/min, the recommendation is 250 to 500 mg every 12 to 24 hours.

ANSWERS:

1. Jane's IBW = 54.7 kg

2. Jane's CrCl = $(140-69)(54.7) / (1.5)(72) = 35.96$ ml/min

NOTE: Since females have 15% less kidney function than males, you must subtract 15% from Jane's CrCl result.

$(0.85)(35.96) = 30.5$ ml/min

3. Since Jane's CrCl (30.5 ml/min) falls within the 10 to 50 ml/min range of the dosage guideline of Keflex, the recommended dose is: 500 mg every 8 to 12 hours.

CREATININE CLEARANCE SAMPLE PROBLEM

NOTE: The CrCl formula will be provided to students during the exam. The emphasis will be on application and making the right assessment.

Joe Smith (JS) is a 72-year old male with a past medical history of COPD. Today he presents to the ED complaining of shortness of breath and wheezing. A respiratory culture from last week reported the presence of *Streptococcus pneumoniae*, consistent with community-acquired pneumonia. Levofloxacin (Levaquin) was ordered, requiring dosage adjustment in a patient with renal insufficiency. Recommend a dosage regimen for levofloxacin (Levaquin), based on Joe's renal function.

A blood chemistry panel reported the following values:

Joe's HT / WT = 5' 9" / 175 LBS.

Na = 148
K = 4.8
Cl = 102
Glucose = 112
CO₂ = 24
BUN = 52
Cr = 1.9

QUESTIONS

1. Calculate Joe's IBW (ideal body wt)
2. Calculate his CrCl (creatinine clearance).
3. Recommend the appropriate dose of levofloxacin for Joe.

Levofloxacin (Levaquin) Monogram

For patients with a CrCl greater than 50 ml/min, the dosage recommendation is levofloxacin is 750 mg IV/PO daily. For patients with CrCl 20-49 ml/min, levofloxacin 750 mg IV/PO every other day (i.e., every 48 hours) is recommended. Patients with a CrCl 10-19 ml/min require a dosage adjustment of levofloxacin 500 mg every other day.

ANSWERS:

1. Joe's IBW = 70.7 kg

2. Joe's CrCl = $(140-72)(70.7) / (1.9)(72) = 35.14$ ml/min

Note: Since Joe is a male patient, we will NOT take into account a 15% reduction in his calculated CrCl value. Female patients would require us to multiply the CrCl result by 0.85 .

3. Since Joe's CrCl value of 35.14 ml/min falls within the 20 to 49 ml/min range in the drug monogram, the recommended dose of levofloxacin is 750 mg IV/ PO every other day (i.e. every 48 hours).